

[Click here to view linked References](#)

Sergio E. Favero-Longo^{1,*}, Enrica Matteucci¹, Paolo Giordani², Alexander G. Paukov³,
Nishanta Rajakaruna^{4,5}

**Diversity and functional traits of lichens in ultramafic areas: a literature-based
worldwide analysis integrated by field data at the regional scale**

¹ Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università di Torino, Viale
Mattioli 25, 10125, Torino, Italy

² Dipartimento di Farmacia, Università di Genova, Viale Cembrano 4, 16148, Genova, Italia
10129, Italy

³ Department of Biodiversity and Bioecology, Institute of Natural Sciences, Ural federal
university, 620000, Ekaterinburg, Russia

⁴ Biological Sciences Department, California Polytechnic State University, San Luis Obispo,
CA 93407, USA

⁵Unit for Environmental Sciences and Management, North-West University, Private Bag
X6001, Potchefstroom, 2520, South Africa

*Corresponding author:

Sergio E. Favero-Longo, PhD.

Università degli Studi di Torino

Dipartimento di Scienze della Vita e Biologia dei Sistemi

Viale Mattioli 25, 10125 Torino, Italy

Tel. +390116705972

Fax +390116705962

sergio.favero@unito.it

Abstract

While higher plant communities found on ultramafics are known to display peculiar characteristics, the distinguishability of any peculiarity in lichen communities is still a matter of contention. Other biotic or abiotic factors, rather than substrate chemistry, may contribute to differences in species composition reported for lichens on adjacent ultramafic and non-ultramafic areas.

This work examines the lichen biota of ultramafics, at global and regional scales, with reference to species-specific functional traits. An updated world list of lichens on ultramafic substrates was analyzed to verify potential relationships between diversity and functional traits of lichens in different Köppen-Geiger climate zones. Moreover, a survey of diversity and functional traits in saxicolous communities on ultramafic and non-ultramafic substrates was conducted in Valle d'Aosta (NW Italy) to verify whether a relationship can be detected between substrate and functional traits that cannot be explained by other environmental factors related to altitude.

Analyses (UPGMA, CCA, SDR) of global lichen diversity on ultramafic substrates (2314 reports of 881 taxa from 43 areas) displayed a zonal species distribution in different climate zones rather than an azonal distribution driven by the shared substrate. Accordingly, variations in the frequency of functional attributes reflected reported adaptations to the climate conditions of the different geographic areas. At the regional scale, higher similarity and lower species replacement were detected at each altitude, independent from the substrate, suggesting that altitude-related climate factors prevail over putative substrate-factors in driving community assemblages. In conclusion, data do not reveal peculiarities in lichen diversity or the frequency of functional traits in ultramafic areas.

Keywords: functional traits; lichen; Köppen-Geiger climate zones; serpentine; ultramafic

Introduction

Deficiency in plant essential nutrients, reduced water availability, combined with Mg:Ca imbalance and often high Ni, Cr and Co — altogether known as the “serpentine factors”— result in unique phanerogamic floras in ultramafic ecosystems (Harrison and Rajakaruna 2011). Ultramafic areas are generally distinct from surrounding non-ultramafic ones in that they harbor low vascular plant diversity and density, leading to sparsely vegetated landscapes (Brooks 1987, with refs therein). Moreover, ultramafic ecosystems display a spectacular richness of plant endemism, species showing exceptional capacity for metal accumulation (i.e. hyperaccumulators; van der Ent et al. 2013), ecotypes with peculiar morphologies (serpentinomorphoses, *sensu* Pichi-Sermolli 1948), abundance of species characterized by disjunct distributions (Rajakaruna 2017), and the co-presence of acidophytic and basiphytic species (e.g. O’Dell and Rajakaruna 2011; van der Ent et al. 2013; Anacker 2014). These features have long fascinated botanists, making ultramafic environments a model system to explore plant adaptation and evolution by natural selection (Harrison and Rajakaruna 2011).

Not surprisingly, investigations on lichens in ultramafic areas have often been discussed with reference to the well-documented peculiarities of higher plant communities on ultramafics (Favero-Longo et al. 2004). However, the recognizability of any peculiarity in lichen communities on ultramafics is still a matter of contention (Favero-Longo 2014). Reduced lichen diversity and abundance have been recognized, but only on certain ultramafic lithologies (Favero-Longo et al. 2015, with refs therein). Many lichen species previously regarded as exclusive to serpentinized ultramafic rocks were subsequently reported from other (mafic) rocks (Favero-Longo et al. 2004). Observations of ‘serpentinomorphoses’ such as stenophyllism or dwarfism were occasionally described in the first half of the last century (Suza 1927; Sambo 1937), but have not been subsequently confirmed. Nevertheless, when lichen diversity has been compared between adjacent ultramafic and non-ultramafic areas, some differences have often been noted, suggesting some substrate-related peculiarities (Favero-Longo et al. 2004; Favero-Longo 2014, with refs therein). Recent work has suggested that other biotic or abiotic factors, rather than rock or soil chemistry, may primarily drive species- and community-level processes in the case of lichens, potentially explaining these peculiarities (Favero-Longo & Piervittori 2009; Rajakaruna et al. 2012; Favero-Longo et al. 2015). However, no investigations have focused on this topic because of the difficulties in comparing studies from different ultramafic areas in different parts of the world.

In this context, the study of functional traits of species (*sensu* Violle et al. 2007) and their direct links to environmental factors has been shown to allow comparisons among different ecosystems and across regions (Giordani et al. 2012). In the case of lichens, the study of morpho-physio-phenological traits influencing growth, reproduction and survival have clarified relationships between lichen traits and environmental factors, including disturbance, solar radiation, water drainage, fire, land management, and climate gradients (Giordani et al. 2014; Nelson et al. 2015; Giordani et al. 2016). Similar analyses of functional traits of lichens with respect to substrate and other macro- and micro-environmental factors have not been performed with respect to lichen communities in ultramafic areas alone or in comparison with those on non-ultramafic substrates. Only some correlations of substrate-related elemental concentrations in thalli with fecundity and secondary metabolite patterns have been suggested for lichens of the genus *Xanthopamelia* on felsic and mafic lithologies (Deduke et al. 2016; Matteucci et al. 2017).

In this paper, we examine the lichen biota in ultramafic areas with reference to species-specific functional traits. Our first objective was to survey diversity and functional traits of lichens in ultramafic areas throughout the world, evaluating the occurrence of local trends of adaptation to geographically-related environmental pressures or of common, ultramafic-related fitness signatures. With this regard, we updated the checklist of lichen reports from ultramafic areas (with respect to Favero-Longo et al. 2004), classified the ultramafic areas — located in different climate regions based on the updated World Map of the Köppen-Geiger climate classification (Kottek et al. 2006; Rubel et al. 2017)— on the basis of the presence or absence of lichen species, and examined whether there is a correlation between lichen diversity and functional traits with their distribution across climate regions. Our second objective was to examine whether a relationship can be detected between substrate and functional traits which cannot be explained by other environmental factors. This work was conducted at a regional scale due to the difficulty in expanding a similar investigation to a wider spatial scale. In particular, diversity and functional traits were surveyed in lichen communities on ultramafic and non-ultramafic substrates in areas of Valle d'Aosta (NW-Italy) at different altitudes, evaluating whether a primary influence of lithology can be detected or whether other environmental factors related to altitude prevail in driving community-level processes at a regional scale.

Methods

World literature survey

A survey was conducted of published records of lichens on ultramafic rocks of stratiform, concentrically zoned, ophiolitic, and high temperature peridotite complexes (see Malpas 1992) throughout the world. Forty-nine pertinent publications were utilized (with some references therein), including floristic and vegetation studies and species reports for 43 ultramafic areas. For each area, information was collected on the ultramafic lithology (e.g. non-serpentinized or serpentinized peridotite) and the climate (according to the updated world map of Köppen-Geiger climate classification; Kottek et al. 2006; Rubel et al. 2017) (Table 1). A comprehensive list of lichen species reported from ultramafic areas was compiled, with nomenclature updated according to Index Fungorum (2017), CNALH (2017) and Nimis (2016). The α -diversity per area and species frequency in the overall areas were analysed. Areas with at least 20 species ($n=26$ areas) were classified (UPGMA, Phi as dissimilarity coefficient, arbitrary resolution of ties; Podani 2001) on the basis of the presence/absence of species with at least two reports. The relative importance of components of γ -diversity [i.e. similarity (S), relativized richness difference (D), and relativized species replacement (R)] was evaluated for all combinations of these areas by analysing the matrix of species presence/absence with SDR Simplex software (2001) using the Simplex method (SDR Simplex; Podani and Schmera 2011). Similarity (S) was calculated following the Jaccard coefficient of similarity:

$$S_{Jac} = a/n$$

where a is the number of species shared by the two plots, and n is the total number of species. The relativised richness difference (D) was calculated as the ratio of the absolute difference between the species numbers of each site (b, c) and the total number of species, n :

$$D = |b-c|/n$$

Relativised species replacement (R) was calculated as:

$$R = 2 * \min \{b, c\}/n$$

A relativised β -diversity as the sum of R+D, a relativised richness agreement as the sum of R+S, and a relativised nestedness as the sum of S+D were also calculated for each pair of areas following Podani and Schmera (2011).

Regional field survey: study area, sampling design and statistical analyses

Lichen diversity was surveyed on serpentinized-ultramafic rocks and non-ultramafic lithologies, including granite, gneiss and micaschist, in nine areas of Valle d'Aosta (NW-Italy) distributed at three different altitudes (approx. 550, 1600 and 2250 m; Table 2). The

intra-alpine Valle d'Aosta (approx. 3200 km²) displays a fairly dry semi-continental climate, ranging from arid (BSk), to temperate (Cfa, Cfb) and boreal (Dfc)-alpine (ET, EF) at the three surveyed altitudes (Rubel et al. 2017a, b).

At each site, delimited on the basis of the homogeneous occurrence of a lithology at a certain altitude, four plots were defined by randomly drawing geographical coordinates determined by a GPS (Garmin 12) and surveyed during the summer of 2012 and 2013. In each plot, three independent 50×50 cm sub-plots were established on the three rock surfaces closest to the randomly extracted coordinate and sharing the following (micro-)environmental features: direct solar irradiation, regular micromorphology, slope <30°, absence of cracks, soil deposits and mosses. Each plot was surveyed using a square grid divided into 25 quadrats (10×10 cm). The frequency of lichen species within each sub-plot (as the sum of their occurrences within the grid quadrats) was estimated visually. Lichens were identified using Clauzade and Roux (1985), Wirth (1995), Smith et al. (2009) and monographic descriptions. Nomenclature follows Nimis (2016). Sample vouchers were deposited at HB-TO *Cryptogamia*.

Areas, plots and sub-plots were classified (UPGMA, Euclidean as dissimilarity coefficient, arbitrary resolution of ties; Podani 2001) on the basis of the frequency of species. The sub-plot level matrix of species frequency was analysed with SDR Simplex software (2001) using the Simplex method (SDR Simplex; Podani and Schmera 2011), as previously detailed.

Functional traits and statistical analyses

For each lichen species listed from at least five ultramafic areas throughout the world (i.e. species reported in >20% of sites with at least 20 species), and for each species listed from ultramafic and non-ultramafic sites of Valle d'Aosta, we defined a set of functional traits (i.e. components of their phenotype that determine their effects on biological processes and their response to environmental factors) (Violle et al. 2007). Following Giordani et al. (2016, with refs therein), we selected traits associated with reproduction and ecophysiology (Table 3). Several traits play a decisive role in the phases of dispersal and establishment of new thalli: most of them include attributes related to morphological characteristics of the spores, such as their shape, number, dimension and color (Armstrong 1981; Morando et al. 2017). Some ecophysiological functions, such as substrate colonization, evapo-transpiration or photon absorption, are strictly related to lichen growth form (Palmqvist 2000), whereas some features (e.g. thallus and epithecium colors or the presence of pruina) are involved in the protection from solar radiation (Giordani et al. 2003). In this regard, secondary metabolites also play protective roles from negative effects of solar radiation, but are also used to regulate pH and

metal homeostasis and as a defense from other abiotic and biotic stresses (Elix & Stocker-Wörgötter 2008; Hauck et al. 2009, 2013). Functional attributes were assigned to each trait (i.e. values or modalities taken by the trait and varying along environmental gradients and/or through time; Giordani et al. 2016). A multidimensional functional space was identified for both the world and regional datasets (lichens found on ultramafics and lichens found on both ultramafic and non-ultramafic substrates, respectively) placing each taxon according to its functional niche and calculating functional distances between species in each dataset.

In particular, the matrices of species presence/absence (world dataset) or frequency (regional dataset, at the plot and sub-plot level level), and those of functional traits were processed through a canonical correspondence analysis (CCA), which partitions variation explained by each variable and constructs a model of significant variables (CCA using biplot scaling for inter-species distances, Hill's scaling for inter-sample distances; choosing forward selection of variables option; performing Monte Carlo permutation test on the first and all ordination axes) (Ter Braak & Verdonschot 1995). The ordinations were performed using CANOCO 4.5 (Ter Braak & Šmilauer 2002).

Results

World survey

A total of 2314 reports of lichens, attributable to 881 specific and subspecific taxa, were listed for ultramafic substrates worldwide (Table S1 in Supplementary Materials). Some tens of other reports, revealing taxonomic uncertainty because of incomplete information (e.g. absence of authority), were excluded from the counts. Most of the taxa were only reported in one (52%) or two (20%) ultramafic areas, while 15% and 4% of taxa in at least five and ten areas, respectively (Fig. 1a). *Candelariella vitellina* was the most commonly reported species on ultramafic substrates, being listed in 28 out of the 43 considered areas. The highest specific richness was reported for the Lizard Peninsula (217 species, Gilbert and James 1987; Fig. 1B), but the different, and not always specified, extent of the surveyed ultramafic areas, together with the heterogeneity of the surveying approaches, may not allow a consistent identification of biodiversity hotspots.

Twenty-six areas, distributed between Europe (up to Urals; 81%) and North (N-) America (19%), hosted more than 20 species. The classification of these areas on the basis of species presence/absence data resulted in four main groups (I-IV; Fig. 2). Group I included areas (n=4) in western coasts of Europe and N-America with warm temperate climate, fully humid

and with a warm summer (Cfb, or Csb bordering on Cfb in the case of Fi). Group II included areas (n=9) with Cfb climate in central Europe, and areas of northern Europe and northern N-America with snow climates, fully humid and with a warm summer (Dfb) or with cold summer and cold winter (Dfc). Group III included (sub-) Mediterranean areas of Europe with warm temperate climate with dry, hot summer (Csa) or more humid areas with hot to warm summer (Cfa-Cfb borders). Group IV included areas of the Alps (n=6), with climates ranging from warm temperate with hot summer, at the border with the Po Plain, to snow to polar climate at highest altitudes (from Cfa-Cfb to ET). Areas of California (n=2) with warm temperate climate with dry and hot summer (Csb) also clustered within this group.

The SDR analysis (Table 4; Fig. 3a) showed a very low species similarity (S), whereas the species replacement (R) was the major component of γ -diversity. Relativized β -diversity (R+D) was approximately 90%.

A number of functional attributes largely characterized the lichen species more widely reported from ultramafic areas (i.e. reported from at least five ultramafic areas; Table 5): presence of chlorococcoid photobiont, a crustose continuous thallus without pruina, predominance of sexual reproduction, and asci with 8 ellipsoid hyaline spores without septa. Such attributes were dominant through all the groups of ultramafic areas I-IV; however, each group had remarkable relative variations in the frequency of functional attributes (see ΔGr in Table 5), as also displayed by CCA-I (Fig.4). The analysis of the world dataset of species presence/absence and the related functional traits of species extracted four axes which accounted for 66.1% of ultramafic area-functional trait relationships. All canonical axes were significant (Monte Carlo test, p-value = 0.002). The first axis (34.9% of correlation) was characterized by growth form (GROW, weighted correlation 0.71), which was the factor exhibiting the higher conditional effect according to forward selection (F-value 6.58, p-value 0.002) and reproductive strategy (REPR, w.c. 0.62, F-value 2.46, p-value 0.002). The production of aliphatic compounds (Alip, F-value 2.44, p-value 0.002), occurrence of pruina (PRUI, F-value 1.87, p-value 0.016) and spore shape (SSHA, F-value 2.17, p-value 0.006) also showed significant conditional effects, being mostly related to axes 2 (12.0% of correlation), 3 (10.7%) and 4 (8.5%), respectively. Ultramafic areas of groups II, III and IV (see Fig. 2) scattered separately along the first axis, while those of group I were separated along the second axis. Group II was positively correlated with GROW and REPR, with reference to higher occurrence of macrolichens (foliose and fruticose) and species with predominant asexual reproduction. Groups III and IV showed no correlation and a negative

correlation, respectively, with GROW and REPR, whereas they were positively correlated to Alip and PRUI, indicating the production of aliphatic compounds and pruina on the thallus surface. Group I was instead negatively or poorly correlated with both GROW and REPR and Alip and PRUI.

Regional survey

A total of 111 lichen taxa were recorded through the nine areas surveyed in Valle d'Aosta, α -diversity per area ranging from 25 to 53 species (Table S4 in Supplementary Materials). Species diversity was not significantly different between ultramafic and non-ultramafic substrates or between different altitudes when evaluated either per area or per plot (ANOVA, $p > 0.005$), although the highest numbers of species characterized were from non-ultramafic areas and plots at medium altitudes (Table 2; Fig. 5). On all substrates, *Candelariella vitellina* was the most commonly reported species in all low altitude areas, while *Rhizocarpon geographicum* was the most commonly reported species at medium and high altitudes. Only these two species, together with *Circinaria caesiocinerea*, occurred in all of the surveyed areas. Species exclusive of ultramafic and non-ultramafic areas were 14% and 44%, respectively. Species occurring at all three altitudes were 16%, while 14%, 24% and 11% were exclusive to areas at low, medium and high altitudes, respectively. The classification of the nine areas on the basis of lichen frequency data resulted in the separation of three main groups, each including ultramafic and non-ultramafic areas, and mostly reflecting the three surveyed altitudes (Fig. 6). A similar pattern was also obtained by analysing the matrix at the plot and sub-plot level (not shown).

The SDR analysis showed a strongly higher similarity (S) for ultramafic and non-ultramafic areas surveyed at the regional scale than that calculated for the world survey limited to ultramafic areas (Table 4; Fig. 3b). Richness difference (D) was instead lower, while species replacement (R) was analogous. Similar values were obtained when the SDR analysis was performed separately for non-ultramafic and ultramafic areas, the latter showing only a slight increase in similarity and decrease in richness difference (Table 4; Fig. 3c, d). Instead, the SDR analysis performed separately for the three altitudes showed remarkably higher similarity and lower species replacement in the case of areas at medium and high altitudes (Table 4; Fig. 3e, f, g).

The functional attributes that dominated the species listed from ultramafic areas at the global scale were also dominant in the lichen species documented at the regional level in ultramafic and non-ultramafic areas of Valle d'Aosta. The frequencies of functional attributes detected at

the regional level, in particular, were close to values calculated for group IV of ultramafic areas, including alpine areas (Table 5). Higher variations in the frequency of attributes (including vegetative, reproductive and metabolic traits) were observed among species from different altitudes relative to species from different substrates (see ΔU -NU and Δalt in Table 5), as also indicated by CCA-II (Fig. 7). The analysis of regional species frequency and related functional traits extracted four axes which only accounted for 64.8% of area-functional trait relationships. All canonical axes were significant (Monte Carlo test, p-value = 0.002). The first axis (35.6% of correlation) was largely characterized by the production of depsides (Deps, weighted correlation 0.44) and aliphatic compounds (Alip, w.c. 0.36) and the type of ascocarp (ATYP, w.c. 0.37) and was negatively correlated with the occurrence of pruina (PRUI, w.c. -0.47), which were all factors exhibiting a high conditional effect according to forward selection (PRUI: F-value 3.04, p-value 0.002; Deps: 2.29, 0.004; Alip 1.83, 0.010; ATYP 1.58, 0.040). The number of spore septa (SSEP, F-value 1.85, p-value 0.020), the number of spores per ascus (SNUM, F-value 1.76, p-value 0.018) and the production of depsidones (Dops, F-value 1.72, p-value 0.022) related to axes 2 (13.8% of correlation), 3 (7.8%) and 4 (7.6%), respectively, and also showed significant conditional effects. Plots from areas at different altitudes scattered separately along the first axis, independent of the substrate. Plots from high altitudes were positively related with Deps, Alip and ATYP, whereas those of low altitudes were positively correlated with PRUI and, subordinately, GROW, which did not show a significant conditional effect.

Discussion

Lichen diversity in ultramafic areas

The number of lichen species worldwide is estimated to be about 25,000-28,000 taxa, but there are less than 15,000 described species (Zedda & Rambold 2015; Scheidegger 2016). Our literature survey shows that more than 5% of these species have been reported on rocks and soils of ultramafic areas, which represent less than 1% of the land surface of Earth (Brooks 1987). Such species percentage may be even higher when considering the many lichenologically unexplored ultramafic areas (as suggested by works in progress in South Africa, Sri Lanka, Portugal, Massachusetts; see Table S7 in Supplementary materials), and that the listed diversity consists of saxicolous and terricolous (s.l.) taxa only directly related to the substrate, while the reported worldwide lichen diversity also includes a wide epiphytic component. Even if we had updated calculations for the overall saxicolous and terricolous lichens, the relevance of lichen diversity in ultramafic areas may not be comparatively

evaluated due to the absence of other world- or continental-scale checklists based on substrate lithology.

The pool of species most frequently reported on ultramafic substrates (4% in more than 10 areas) includes common, widely-distributed species often reported from silicate-rocks and related soils worldwide (see Wirth 1972). By contrast, the majority of species, which drive the overall diversity, were only reported from one or two areas. SDR analysis showed low similarity ($S=10.5$) across ultramafic areas, which, together with a high replacement ($R=55.5$), determines a high anti-nestedness ($S+R=66.0$), generally indicating a zonal species distribution with high species turnover (Podani & Schmera 2011), rather than an azonal distribution driven by the shared substrate. This pattern agrees with the paucity of endemic lichen species reported from ultramafics, in contrast to the high endemism of phanerogamic communities on ultramafics. Only 8 lichen species, 5 of which are known only from their type localities, can be considered serpentine endemics (Table S8 in Supplementary Materials); however, there is potential that these species may be found on other mafic substrates in the future, as has been the case with other previously reported serpentine-endemic lichen species (Wirth 1972).

The classification of the ultramafic areas on the basis of the presence or absence of lichen species generally reflects their distribution in different climate zones, with reference to the updated Köppen-Geiger classification. Climate (rather than the substrate) may be the primary driver of lichen diversity in ultramafic areas. Although phylogeographic studies have only recently started to address biogeographic histories of lichens (Divakar and Crespo 2015), the recognized groups I-IV and the related sub-groups mostly reflect traditionally distinguished lichen biogeographical domains and their relationships (Seaward 1977; Galloway 2009): the highly humid climates of western Europe and western N-America (group I), the boreal coniferous zone (group II), and Southern Europe (groups III and IV), with a Oromediterranean element remarkably related to the central Europe lichen vegetation (Nimis 1996), an alpine element, and some relationships with the “Mediterranean” coast of California. A substrate-specificity is well known for saxicolous and terricolous lichens and is displayed by strong divergence between communities on silicate and carbonate substrates (Brodo 1973). However, a similar divergence may not be seen when lichen communities on different silicate substrates are compared, including serpentinites. Adaptive strategies for living on silicate (and not on carbonate) substrates may be sufficient to cope with the serpentine-factors (see sections below).

Lichen functional traits in ultramafic areas

A series of functional attributes dominate through all the groups (I-IV) of ultramafic areas, but they merely mirror their dominance in the global lichen biota (e.g. the dominance of chlorococcoid photobionts, crustose species, sexually-reproducing species, asci with 8 simple, ellipsoid, hyaline spores). The more or less noticeable frequency variation of these dominant attributes in favour of other less common attributes (see Table 5) reflects the climate conditions of the different geographic areas and their typical lichen vegetation, independent of the substrate lithology. Highest variations among groups (Δ group max.), recorded for the growth form of thalli, are related to the higher frequency of crustose species in alpine areas, and in the arid areas of California (group IV), than in the more humid areas of the boreal (group II) or Oromediterranean regions of Europe (group III), richer for terricolous fruticose species, while foliose taxa prevail in the oceanic western coasts of Europe (group I) (Seaward 1977). The higher prevalence of sexual reproductive strategies in the alpine areas (group IV) agrees with the prevalence of sexual reproduction in lichens of extreme habitats (Seymour et al. 2005). A higher pigmentation of epithecium (and spores) also characterizes the alpine areas (group IV), playing a protective role against excessive radiation (e.g. Elix & Stocker-Wörgötter 2008; Nguyen et al. 2013). Higher occurrence of pruina and aliphatic compounds in species of group IV also agrees with the necessity of alpine lichens to tolerate stresses related to high radiation and low temperatures (e.g. Giordani et al. 2003; Boustie et al. 2011). Accordingly, in CCA-I, functional traits GROW, REPR, PRUI and Alip are significant conditional factors in driving the divergence among lichen communities of ultramafic areas in different climate regimes.

We can thus suggest the occurrence of local trends of adaptation to geographically related environmental pressures more than recognizing common, ultramafic-related fitness signatures. On the other hand, as in the evaluation of the lichen specific richness through the ultramafic areas, the absence of a similar worldwide survey of functional traits for lichens on non-ultramafic substrates prevents a comparative analysis to recognize shifts of attribute frequencies potentially related to the substrate, which we instead evaluated at the regional level.

Are there peculiarities in lichen communities in ultramafic areas?

The comparative analysis of ultramafic and non-ultramafic areas in the Valle d'Aosta did not show significant paucity of species on ultramafics, in contrast to the paucity of phanerogams commonly reported from ultramafics (Favero-Longo 2014). Dominant species were the same

in both ultramafic and non-ultramafic areas, in agreement with their prevalence on silicate (s.l.) areas in Europe and worldwide (see e.g. Wirth 1972). Some species were exclusively recorded in ultramafic or non-ultramafic areas, as previously reported in comparative analysis of serpentine and adjacent mafic outcrops (e.g. Sirois 1987; Paukov and Trapeznikova 2005; Favero-Longo & Piervittori 2009; Paukov 2009; Rajakaruna et al. 2012) and similarity (S) in SDR analysis was slightly higher within ultramafic than in non-ultramafic areas, suggesting the possibility of some substrate-related species distributions. However, the species replacement (R) was the same for ultramafic and non-ultramafic areas, and the lower similarity in non-ultramafic areas seems mostly related to differences in diversity richness rather than to peculiar species occurrences. More remarkably, higher similarity and lower species replacement detected at each altitude, independent from the substrate (SDR in Table 4, and UPGMA classification in Fig. 6), suggest that altitude-related climate factors prevail over putative substrate-factors in driving community assemblages at the regional scale. Accordingly, climate-related factors such as water availability and solar radiation were shown to be the dominant drivers of the probability of lichen occurrence at the micro-scale (Giordani et al. 2014).

As expected, functional attributes characterizing the surveyed ultramafic areas in the Valle d'Aosta agree with the trend shown at the world-scale by areas of group IV, including alpine areas: high presence of crustose species, high frequency of sexual reproduction, occurrence of pruina, but also similar patterns of frequency for attributes assigned to thallus continuity/discontinuity and color, and reproductive traits. However, the same (and even higher) agreement was recognized when the surveyed areas were considered altogether and also for the non-ultramafic areas themselves. Moreover, the variation in the frequency of functional attributes was more pronounced between the areas at the different altitudes than between ultramafic and non-ultramafic areas, confirming the primary dependence of lichen assemblages on climate-related factors (varying at different altitudes) than on substrate features. This agrees with the previously reported relationships of lichens with atmospheric-related factors such as humidity (water supply), temperature (dry-wet cycles, metabolic activity), solar radiation (photosynthetic efficiency) (Nash 2008), and a subordinate role of the “substrate compartment”, despite its involvement in mineral nutrition (Favero-Longo 2014 with refs therein).

On the other hand, the fact that saxicolous lichen communities with similar functional attributes colonize all silicate substrates, without any remarkable peculiarity for ultramafic areas, suggests that lichen adaptation to environmental pressures related to silicate rocks also

satisfy the tolerance of “serpentine factors”, including low nutrient availability and high heavy metal concentrations. Low nutrients, by limiting the phanerogamic component, may be considered a positive factor for lichens, favouring their presence due to lower competition (Favero-Longo & Piervittori 2009). Many studies have considered lichen responses to heavy metals, showing remarkable advances of knowledge with regard to the role of secondary metabolites in modulating pH and metal homeostasis (Hauck et al. 2009, 2013). For example, variation in the production of secondary metabolites with differential affinity to iron was shown to regulate the presence or absence of certain species on iron-rich substrates (Hauck et al. 2007). Similar metabolic features may be related to the wide-spectrum adaptation of saxicolous lichens to ultramafic and non-ultramafic silicate substrates (Favero-Longo et al. 2015), but the analysis of this point goes beyond the aims of this paper and would need direct testing on the contents of secondary metabolites in species (and even thalli) from the surveyed plots (Matteucci et al. 2017). However, on the basis of literature on the specific production of secondary metabolites, the regional survey conducted here shows differences in the frequency of species producing metabolites of different classes (e.g. depsidones) both between ultramafic and non-ultramafic areas and at different altitudes. This finding suggests the need to examine the production of secondary metabolites potentially involved in metal homeostasis not only on different substrates, but also under different climate conditions, which may affect mineral cycling and element availability.

In conclusion, rather than searching for peculiarities in diversity and functional traits, lichenological research may focus on ultramafic environments as natural labs (see Harrison and Rajakaruna 2011) to understand adaptation to metal stresses and the potential role of secondary metabolites in metal homeostasis.

Acknowledgements

EM was the recipient of a postdoctoral fellowship (RICERCA FSE 2011-2012, funded by the European Union, Italian Ministry of Labour and Social Policies, Regione Autonoma Valle d’Aosta: agreement 10016/DPLF/28June2012). The authors are grateful to Bianca Bovero (University of Torino) for her assistance during fieldwork and to Rosanna Piervittori (University of Torino) for helpful discussions. The authors thank two reviewers for their useful and constructive comments which helped improve the quality of the manuscript. The work of AP is financially supported by RFBR (grants 15-04-05971 and 16-04-01346).

References

- Anacker BL (2014). The nature of serpentine endemism. *Amer. J. Bot.* 10:219-224. doi: 10.3732/ajb.1300349
- Aptroot A, John V (2015) An historical lichen collection from New Caledonia. *Herzogia* 28:307-321. doi:10.13158/heia.28.2.2015.307
- Aptroot A, Van de Vijver B, Lebouvier M, Ertz D (2011) Lichens of Ile Amsterdam and Ile Saint Paul (TAAF, southern Indian Ocean). *Nova Hedwigia* 92:343-367. doi:10.1127/0029-5035/2011/0092-0343
- Armstrong RA (1981). Field experiments on the dispersal, establishment and colonization of lichens on a slate rock surface. *Environ. Exp. Bot.* 21:115-120. doi: 10.1016/0098-8472(81)90016-2
- Bilovitz PO, Mayrhofer H (2009) A contribution to the lichenized fungi of Bosnia and Herzegovina. *Fritschiana* (Graz) 65:53-56.
- Boustie J, Tomasi S, Grube M (2011) Bioactive lichen metabolites: alpine habitats as an untapped source. *Phytochem. Rev.* 10:287-307. doi:10.1007/s11101-010-9201-1
- Brodo IM (1973) Substrate ecology. In: Ahmadjian V, Hale ME (eds) *The lichens*. Academic Press, New York, 401-441
- Brooks RR (1987) *Serpentine and its vegetation: a multidisciplinary approach*. Dioscorides Press, Croom Helm.
- Candan M, Türk AO (2008). Lichens of Malatya, Elazığ and Adıyaman provinces (Turkey). *Mycotaxon* 105:19-22.
- Clauzade G, Roux C (1985) *Likenoj de Okcidenta Europo. Ilustrita determinlibro*. Bull. Soc. Bot. Centre-Ouest 7:3-893.
- CNALH (accessed 9 September 2017) available at <http://lichenportal.org/portal/>
- Deduke C, Halden NM, Piercey-Normore MD (2016) Comparing element composition of rock substratum with lichen communities and the fecundity of *Arctoparmelia* and *Xanthoparmelia* species. *Botany* 94:41-51. doi: 10.1139/cjb-2015-0141
- Divakar PK, Crespo A (2015) Molecular phylogenetic and phylogenomic approaches in studies of lichen systematics and evolution. In: Upreti DK, Divakar PK, Shukla V, Bajpai R (eds) *Recent advances in lichenology. Modern methods and approaches in lichen systematics and culture techniques, Volume 2*. Springer, New Delhi, pp 45-60

493 Elix JA, Stocker-Wörgötter E (2008) Biochemistry and secondary metabolites. In: Nash III
 494 TH (ed) Lichen biology. Cambridge University Press, Cambridge, pp 104-133

495 Favero-Longo SE, Piervittori R (2009). Measuring the biodiversity of saxicolous lichens
 496 above timberline with reference to environmental factors: the case-study of a Natura 2000
 497 site of western Alps. *Phytocoenologia* 39:51-78. doi:10.1127/0340-269X/2009/0039-0051

498 Favero-Longo SE, Siniscalco C, Piervittori R (2006). Plant and lichen colonization in an
 499 asbestos mine: spontaneous bioattenuation limits air dispersion of fibres. *Pl. Biosyst.*
 500 140:190-205. doi: 10.1080/11263500600756546

501 Favero-Longo SE, Matteucci E, Morando M, Rolfo F, Harris T, Piervittori R (2015). Metals
 502 and secondary metabolites in saxicolous lichen communities on ultramafic and non-
 503 ultramafic rocks of the Western Italian Alps. *Austral. J. Bot.* 63:276-291. doi:
 504 10.1071/BT14256

505 Favero-Longo SE, Isocrone D, Piervittori R (2004). Lichens and ultramafic rocks: a review.
 506 *Lichenologist* 36:391-404. doi:10.1017/S0024282904014215

507 Favero-Longo SE (2014) Lichens on metal-rich substrates. In: Rajakaruna N, Boyd R, Harris
 508 T (eds) Plant ecology and evolution in harsh environments. Nova Publishers, New York,
 509 pp 53-76.

510 Fryday AM (2005). The genus *Porpidia* in northern and western Europe, with special
 511 emphasis on collections from the British Isles. *Lichenologist* 37:1-35. doi:
 512 10.1017/S0024282904014628

513 Gallo LM, Piervittori R (1991) La flora lichenica rupicola dei Monti Pelati di Baldissero
 514 (Canavese, Piemonte). In: Giachino PM (ed) I Monti Pelati di Baldissero. Importanza
 515 paesistica e scientifica. Atti Convegno 1991, pp. 25-31.

516 Galloway DJ (2009) Lichen biogeography. In: Nash III TH (ed) Lichen biology. Cambridge
 517 University Press, Cambridge, pp 315-335

518 Gilbert OL, James PW (1987) Field meeting on the Lizard Peninsula, Cornwall. *Lichenologist*
 519 19:319-334. doi:10.1017/S0024282987000288

520 Gilbert OL (1983) The lichens of Rhum. *Trans. Bot. Soc. Edinburgh* 44:139-149.
 521 doi:10.1080/03746608308685380

522 Giordani P, Benesperi R, Rizzi G, Brunialti G (2009). New records for lichen regional floras
 523 of Italy. *Webbia* 64:153-158. doi:10.1080/00837792.2009.10670855

524 Giordani P, Modenesi P, Tretiach M (2003). Determinant factors for the formation of the
 525 calcium oxalate minerals, weddellite and whewellite, on the surface of foliose lichens.
 526 *Lichenologist* 35:255-270. doi: 10.1016/S0024-2829(03)00028-8

527 Giordani P, Brunialti G, Bacaro G, Nascimbene J (2012). Functional traits of epiphytic
 528 lichens as potential indicators of environmental conditions in forest ecosystems. *Ecol.*
 529 *Indicators* 18:413-420. doi:10.1016/j.ecolind.2011.12.006

530 Giordani P, Incerti G, Rizzi G, Rellini I, Nimis PL, Modenesi P (2014) Functional traits of
 531 cryptogams in Mediterranean ecosystems are driven by water, light and substrate
 532 interactions. *J. Veg. Sci.* 25:778-792. doi:10.1111/jvs.12119

533 Giordani P, Rizzi G, Caselli A, Modenesi P, Malaspina P, Mariotti MG (2016) Fire affects the
 534 functional diversity of epilithic lichen communities. *Fungal Ecology* 20:49-55.
 535 doi:10.1016/j.funeco.2015.11.003

536 Hafellner J (1991) Die Flechtenflora eines hochgelegenen Serpentinistockes in den Östalten
 537 (Österreich, Steiermark). *Mitt. Naturwiss. Vereins Steiermark, Graz* 121:95-106.

538 Hafellner J (2001) Bemerkenswerte Flechtenfunde in Österreich. *Fritschiana* 28:1-30.

539 Hakulinen R (1958) Jäkälä Suomen serpentiinikallioilta. *Ann. Bot. Soc. Zool.-Bot. Fenn.*
 540 "Vanamo" 12:143-145.

541 Hansen CJ, Goertzen LR (2012). *Psora icterica* (Lecanorales, Psoraceae), a new and
 542 interesting disjunction from Alabama. *Opuscula Philolichenum* 11:49-51.

543 Harris TB, Olday FC, Rajakaruna N. (2007) Lichens of Pine Hill, a peridotite outcrop in
 544 eastern North America. *Rhodora* 109:430-447. doi: 10.3119/0035-
 545 4902(2007)109[430:LOPHAP]2.0.CO;2

546 Harrison S, Rajakaruna N (2011) *Serpentine. The evolution and ecology of a model system.*
 547 University of California Press, Berkeley.

548 Hauck M, Huneck S, Elix JA, Paul A (2007). Does secondary chemistry enable lichens to
 549 grow on iron-rich substrates? *Flora* 202:471-478. doi: 10.1016/j.flora.2006.08.007

550 Hauck M, Jürgens S-R, Willenbruch K, Huneck S, Leuschner C (2009) Dissociation and
 551 metal-binding characteristics of yellow substances suggest a relationships with site
 552 preferences of lichens. *Ann. Bot.* 103:13-22. doi:10.1093/aob/mcn202

553 Hauck M, Böning J, Jacob M, Dittrich S, Feussner I, Leuschner C (2013). Lichen substance
 554 concentrations in the lichen *Hypogymnia physodes* are correlated with heavy metal

555 concentrations in the substratum. Environm. Exp. Bot. 85:58-63.
 556 doi:10.1016/j.envexpbot.2012.08.011
 557 Huneck S, Yoshimura I (1996) Identification of lichen substances. Springer, Berlin.
 558 Index Fungorum (accessed 9 September 2017) available at: <http://www.indexfungorum.org/>
 559 Isocrono D, Favero Longo SE, Piervittori R (2008). La flora lichenica del Parco Naturale del
 560 Mont Avic (Valle d'Aosta, Italia). Rev. Valdôtaine Hist. Nat. 62:77-93.
 561 Kantvilas G (1991) Records of east African lichens in cool temperate Australia. Nordic J. Bot.
 562 11:369-373. doi:10.1111/j.1756-1051.1991.tb01416.x
 563 Kocakaya M, Halici MG, Aksoy A (2014) Lichenized and lichenicolous fungi of Gevne
 564 Valley (Konya, Antalya). Turk J Bot 38:358-369. doi:10.3906/bot-1303-29
 565 Kossowska M (2001) Epilithic lichens on serpentinite rocks in Poland. Polish Bot. J. 46:191-
 566 197
 567 Kottek M, Grieser J, Beck C, Rudolf B, Rubel F (2006). World map of the Köppen-Geiger
 568 climate classification updated. Meteorol. Z. 15:259-263. doi:10.1127/0941-
 569 2948/2006/0130
 570 Krause W, Klement O (1958) Zur Kenntnis der Flora und Vegetation auf Serpentinstandorten
 571 des Balkans. 3. Felsflechten-Gesellschaften im Gosrovic-Gebiet (Bosnien) und Zlatibor-
 572 Gebirge (Serbien). Vegetatio 8:1-19.
 573 Krause W., Klement O (1962) Zur Kenntnis der Flora und Vegetation aus Serpentinstandorten
 574 des Balkans. 5. Flechten und Flechtengesellschaften auf Nord-Euboa (Griechenland).
 575 Nova Hedwigia 4: 189-262.
 576 Kretschmer L (1931) Die Pflanzengesellschaften auf Serpentin im Gurhofgraben bei Melk.
 577 Verh. Zool.-Bot. Ges. Wien 80:163-208.
 578 Krzewicka (2009) The 'Verrucaria fuscella group' in Poland with some nomenclatorial
 579 remarks. Acta Soc. Bot. Poloniae 78:229-234. doi:10.5586/asbp.2009.029
 580 Lammermayr L. (1934) Übereinstimmung und Unterschiede in der Pflanzendecke über
 581 Serpentin und Magnesit. Mitteilungen der Naturwissenschaftlichen Vereins für Steiermark,
 582 Graz 71:41-62.

583 Malpas J (1992) Serpentine and the geology of serpentinized rocks. In: Roberts A, Proctor J
584 (eds) The ecology of areas with serpentinized rocks. A world view. Kluwer Academic
585 Publishers, Dordrecht, pp 7-30

586 Matteucci E, Bovero B, Vanacore Falco I, Piervittori R (2015) Comunità licheniche rupicole
587 in Valle d'Aosta: confronti a quote differenti. *Rev. Valdôtaine Hist. Nat.* 69:61-77.

588 Matteucci E., Occhipinti A., Piervittori R., Maffei ME, Favero-Longo SE (2017)
589 Morphological, secondary metabolite and ITS (rDNA) variability within usnic acid-
590 containing lichen thalli of *Xanthoparmelia* explored at the local scale of rock outcrop in W-
591 Alps. *Chemistry and Biodiversity* 14:e1600483. doi:10.1002/cbdv.201600483

592 Medeiros ID, Fryday AM, Rajakaruna N (2014) Additional lichen records and mineralogical
593 data from metal-contaminated sites in Maine. *Rhodora* 116:323-347. doi:10.3119/13-26

594 Morando M, Favero-Longo SE, Carrer M, Matteucci E, Nascimbene J, Sandrone S,
595 Appolonia L, Piervittori R (2017). Dispersal patterns of meiospores shape population
596 spatial structure of saxicolous lichens. *Lichenologist* 49:397-413. doi:
597 10.1017/S0024282917000184

598 Nash III TH (2008) Lichen biology. Cambridge University Press, Cambridge.

599 Nelson PR, McCune B, Roland C, Stehn S (2015). Non-parametric methods reveal non-linear
600 functional trait variation of lichens along environmental and fire age gradients. *J. Veg. Sci.*
601 26: 848-865. doi:10.1111/jvs.12286

602 Nguyen KH, Chollet-Krugler M, Gouault N, Tomasi S (2013). UV-protectant metabolites
603 from lichens and their symbiotic partners. *Nat. Prod. Rep.* 30:1490-1508.
604 doi:10.1039/C3NP70064J

605 Nimis PL (1996) Towards a checklist of Mediterranean lichens. *Bocconea* 6:5-17.

606 Nimis PL (2016). The lichens of Italy. A second annotated catalogue. EUT, Trieste.

607 O'Dell RE, Rajakaruna N (2011) Intraspecific variation, adaptation, and evolution. In:
608 Harrison S, Rajakaruna N (eds) Serpentine. The evolution and ecology of a model system.
609 University of California Press, Berkeley, pp 97-137

610 Palmqvist K. (2000). Tansley review no. 117: carbon economy in lichens. *New Phytol.*
611 148:11-36. doi: 10.1046/j.1469-8137.2000.00732.x.

612 Paukov AG, Trapeznikova SN (2005) Lithophilous lichens of Middle Urals. *Folia Cryptog.*
613 *Estonica* 41:81-88.

- 614 Paukov AG (2009). The lichen flora of serpentine outcrops in the Middle Urals of Russia. N.
615 E. Naturalist 16 (Special Issue No. 5):341-350.
- 616 Paukov A, Sipman HJM, Kukwa M, Repin R, Teptina A (2017) New lichen records from the
617 mountains Kinabalu and Tambuyukon (Kinabalu Park, Malaysian Borneo). Herzogia
618 30:237-252.
- 619 Peksa O (2011) Lišejníky národní přírodní památky Křížky. Sborník muzea Karlovarského
620 kraje 19:259-272.
- 621 Pichi-Sermolli R (1948) Flora e vegetazione delle serpentine e delle altre ofioliti dell'alta
622 valle del Tevere (Toscana). Webbia 6:1-380. doi:10.1080/00837792.1948.10669585
- 623 Podani J (2001) SYN-TAX 2000. Computer Programs for Data Analysis in Ecology and
624 Systematics. User's Manual. Scientia Publishing, Budapest.
- 625 Podani J, Schmera D (2011). A new conceptual and methodological framework for exploring
626 and explaining pattern in presence-absence data. Oikos 120:1625-1638.
627 doi:10.1111/j.1600-0706.2011.19451.x
- 628 Poelt J (1975) *Squamarina serpentinii* species nova (Lichenes, Lecanoraceae) aus Serbien.
629 Herzogia 3:425-432.
- 630 Purvis OW, Halls C (1996) A review of lichens in metal-enriched environments.
631 Lichenologist 28:571-601. doi:10.1006/lich.1996.0052
- 632 Rajakaruna, N. 2017. Lessons on evolution from the study of edaphic specialization. Bot.
633 Rev. 83 (in press). doi:10.1007/s12229-017-9193-2
- 634 Rajakaruna N., Knudsen K., Fryday AM, O'Dell RE, Pope N., Olday FC, Woolhouse S
635 (2012) Investigation of the importance of rock chemistry for saxicolous lichen communities
636 of the New Idria serpentinite mass, San Benito County, California, USA. Lichenologist
637 44:695-714. doi:10.1017/S0024282912000205
- 638 Ritter-Studnička H, Klement O (1968) Über Flechtenarten und deren Gesellschaften auf
639 Serpentin in Bosnien. Oesterr. Bot. Z. 115:93-99.
- 640 Rubel F, Brugger K, Haslinger K, Auer I (2017). The climate of the European Alps: Shift of
641 very high resolution Köppen-Geiger climate zones 1800–2100. Meteorol. Z. 26:115-125.
642 doi:10.1127/metz/2016/0816
- 643 Ryan BD (1988) Marine and maritime lichens on serpentine rocks on Fidalgo Island,
644 Washington. Bryologist 91:186-190. doi:10.2307/3243217

645 Sambo E (1927) I licheni del Monte Ferrato (Toscana). *Nuovo Giornale Botanico Italiano*
646 34:333-358.

647 Sambo E (1937) Sull'azione vicariante del magnesio invece del calcio in licheni calcicoli di
648 roccia alcalica. *Nuovo Giorn. Bot. Ital.* 24:246-250. doi:10.1080/11263503709437956

649 Sánchez-Biezma MJ, López de Silanes ME (1999) *Porpidia nadvornikiana*, a species of
650 ultrabasic rocks: second record for Europe. *Lichenologist* 31:637-639. doi:
651 10.1006/lich.1999.0238

652 Sánchez-Biezma MJ, Carballal R, López de Silanes ME (1996) Algunos líquenes de rocas
653 básicas y ultrabásicas nuevos para el nord de España. *Cryptogamie, Bryologie,*
654 *Lichénologie* 17:203-211.

655 Sánchez-Biezma Serrano MJ, Alvarez Andrés J, López de Silanes Vázquez ME (2001)
656 Líquenes de las rocas ultramáficas en la Sierra de A Capelada (A Coruña, NW de España).
657 *Bot. Complut.* 25:261-269.

658 Scheidegger C (2016) As thick as three in a bed. *Molec. Ecol.* 25:3261-3263.
659 doi:10.1111/mec.13710/full

660 Seaward MRD (1977) *Lichen ecology*. Academic Press, London.

661 Seymour FA, Crittenden PD, Dyer PS (2005). Sex in the extremes: lichen-forming fungi.
662 *Mycologist* 19:51-58. doi:10.1017/S0269-915X(05)00201-6

663 Sigal LL (1989) The lichens of serpentine rocks and soils in California. *Mycotaxon* 34:221-
664 238.

665 Sipman HJM (1993). Lichens from Mount Kinabalu. *Trop. Bryol.* 8:281-314.

666 Sirois L, Lutzoni F, Grandtner MM (1987) Les lichens sur serpentine et amphibolite du
667 plateau du mont Albert, Gaspésie, Québec. *Canad. J. Bot.* 66:851-862. doi: 10.1139/b88-
668 124

669 Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW, Wolseley PA (2009)
670 *The lichens of Great Britain and Ireland*. British Lichen Society, London.

671 Suza J (1927) Lichenologický ráz západočeských serpentinů. *Čas. Morav. Zemsk. Mus.*
672 25:251-281.

673 Takala K, Seaward MRD. (1978) Lichens of the Niinivaara serpentinite region, E. Finland.
674 *Memoranda Soc. Fauna Fl. Fenn.* 54:59-63.

675 Ter Braak CJF, Šmilauer P (2002) CANOCO reference manual and CanoDraw for Windows
676 User's guide: software for canonical community ordination (version 4.5). Microcomputer
677 Power, Ithaca (NY).

678 Ter Braak CJ, Verdonschot PF (1995). Canonical correspondence analysis and related
679 multivariate methods in aquatic ecology. *Aquatic Sci.* 57:255-289.
680 doi:10.1007/BF00877430

681 van der Ent A., Baker AJ, Reeves RD, Pollard AJ, Schat H (2013). Hyperaccumulators of
682 metal and metalloid trace elements: facts and fiction. *Pl. & Soil* 362:319-334.
683 doi:10.1007/s11104-012-1287-3

684 Versegghy K (1974) Nachtrag II. zum "Typenverzeichnis der Flechtensammlung in der
685 Botanischen Abteilung des Ungarischen Naturwissenschaftlichen Museums". *Studia Bot.*
686 Hung. 9:23-29.

687 Vězda A (1972) *Haplocarpon nadvornikianum* Vězda sp. nov. *Preslia* 44:208-212.

688 Violle C, Navas ML, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E (2007) Let the
689 concept of trait be functional! *Oikos* 116:882-892. doi: 10.1111/j.0030-1299.2007.15559.x

690 von Brackel W (2007). Zur Flechtenflora der Serpentinittfelsen in Nordostbayern. *Hoppea* 68,
691 253-268.

692 Wirth V (1972) Die Silikatflechten Gemeinschaften im außeralpinen Zentraleuropa. Diss.
693 Bot. 17:1-306.

694 Wirth V (1995) Die Flechten. Baden-Württembergs. Ulmer, Stuttgart.

695 Zahlbruckner A (1907) Aufzählung der von Dr. H. Bretzl in Griechenland gesammelten
696 Flechten. *Hedwigia* 47:60-65.

697 Zedda L, Rambold G (2015) The diversity of lichenised fungi: Ecosystem functions and
698 ecosystem services. In: Upreti DK, Divakar PK, Shukla V, Bajpai R (eds) Recent advances
699 in lichenology. Modern methods and approaches in lichen systematics and culture
700 techniques, Volume 2. Springer, New Delhi, pp 121-145.

701

Figure captions

Fig. 1 Frequency and diversity of lichen taxa from ultramafic areas worldwide. (a) Number of taxa which were reported from 1 to 28 times from the ultramafic areas considered; (b) number of species listed through the 43 ultramafic areas.

Fig. 2 Classification of ultramafic areas (n=26, with >20 lichen taxa) on the basis of species presence/absence data and their world distribution. (a) UPGMA Classification: group I, diamonds; group II, circles; group III, squares; group IV, boxes. Different symbol colours indicate different sub-groups (not named). Abbreviations for areas are listed in Table 1; information on climates follows Kottek et al. (2006) and Ruben et al. (2017). FL, floristic/vegetation studies; SR, species reports. Cophenetic correlation; 0.82. (b, c), distribution of the classified ultramafic areas through the world (Europe in inset (c)).

Fig. 3 SDR simplex ternary plots for the world (a) and regional (b-f) datasets. Color patterns refer to Kernel densities of site pairs included in the datasets. World level (a): all pairs of ultramafic areas. Regional level: (b) all pairs of sub-plots, (c) all pairs of sub-plots in ultramafic areas, (d) all pairs of sub-plots in non-ultramafic areas, (e-f) all pairs of sub-plots at low, medium and high altitudes, respectively. The abbreviations S, D and R refer to relative similarity, richness difference, and species replacement, respectively.

Fig. 4 Factorial maps in the canonical correspondence analysis (CCA-I) showing (a) the position of ultramafic areas considered at the world scale (abbreviations according to Table 1; symbols according to UPGMA classification in Fig. 2) together with (b) the contributions of different functional traits: photosynthetic strategy (PHOT), growth form (GROW), thallus continuity (TCON), color of the thallus (TCOL), occurrence of pruina (PRUI), reproductive strategy (REPR), ascocarp type (ATYP), color of the epithecium /disc (ACOL), number of ascospores in each ascus (SNUM), spore shape (SSHA), number of septa (SSEP), color (SCOL) and length (SLEN) of the spores, production of secondary metabolites as aliphatic compounds (Alip), quinones (Quin), chromones and xanthonones (Chro), pulvinic derivatives (Pulv), depsides (Deps), depsidones (Dops), dibenzofuranes (Dibe), terpenoids (Terp) (functional traits of each species in Table S2 in Supplementary Materials). All the extracted axes displayed in the figure were significant according to Monte Carlo test. Functional traits exhibiting significant conditional effects are in bold on the diagram (scores in Table S3 in Supplementary Materials).

736

737 **Fig. 5** Lichen richness in ultramafic and non-ultramafic areas of Valle d'Aosta. (a)
738 Localization of surveyed areas in Valle d'Aosta (NW-Italy, as visualized in the left bottom
739 corner) for ultramafic (black symbols) and non-ultramafic (grey symbols) substrates at low
740 (down triangle), medium (circle) and high (up triangle) altitudes (scale bar: 15 km); (b)
741 species diversity per plot in ultramafic and non-ultramafic areas at the different altitudes.
742

743 **Fig. 6** UPGMA classification of the surveyed ultramafic and non-ultramafic areas of Valle
744 d'Aosta on the basis of specific frequency data. Symbols clustered in groups A-C indicate
745 ultramafic (black symbols) and non-ultramafic (grey symbols) areas at low (down triangle),
746 medium (circle) and high (up triangle) altitudes. Cophenetic correlation; 0.90.
747

748 **Fig. 7** Factorial maps in the canonical correspondence analysis (CCA-II) showing the position
749 of ultramafic and non-ultramafic areas surveyed at the regional scale of Valle d'Aosta
750 (symbols according to Figs. 5 and 6) together with contributions of the different functional
751 traits (abbreviations as in Fig. 4B): photosynthetic strategy (PHOT), growth form (GROW),
752 thallus continuity (TCON), color of the thallus (TCOL), occurrence of pruina (PRUI),
753 reproductive strategy (REPR), ascocarp type (ATYP), color of the epithecium /disc (ACOL),
754 number of ascospores in each ascus (SNUM), spore shape (SSHA), number of septa (SSEP),
755 color (SCOL) and length (SLEN) of the spores, production of secondary metabolites as
756 aliphatic compounds (Alip), quinones (Quin), chromones and xanthonones (Chro), pulvinic
757 derivatives (Pulv), depsides (Deps), depsidones (Dops), dibenzofuranes (Dibe), terpenoids
758 (Terp) (functional traits of each species in Table S5 in Supplementary Materials). All the
759 extracted axes displayed in the figure were significant according to Monte Carlo test.
760 Functional traits exhibiting significant conditional effects are in bold on the diagram (scores
761 in Table S6 in Supplementary Materials).

762

763 **Tables**

764 Table 1. Ultramafic areas surveyed with regard to lichen diversity or cited in species reports. Abbreviations of areas with more and less than 20
765 reported lichen taxa are reported out of and in parentheses, respectively. Information on climate according to the updated world map of Köppen-
766 Geiger climate classification (Kottek et al. 2006; Rubel et al. 2017). FL, floristic-vegetation study; SR, species report.

Geographic area				Climate	Ultramafic lithology	Number of lichen taxa	Source Reference	Type
Abbr.	Location	Country	Continent					
Al	Mont Albert, Gaspésie	Canada	N-America	Dfc	Serpentinite	139	Sirois et al. (1987)	FL
Av	Mt. Avic	Italy	Europe	Dfc-ET	Serpentinite	91	Isocrono et al. (2008); Favero-Longo and Piervittori (2009)	FL
Ba	NE-Bavaria	Germany	Europe	Cfb	Serpentinite	129	von Brackel (2007)	FL
Bl	Balangero asbestos mine	Italy	Europe	Cfb-Cfa	Serpentinite	23	Favero-Longo et al. (2006)	FL
(BM)	Mohelno and Raškov, Bohemian-Moravian Highlands	Czech Republic	Europe	Cfb	Serpentinite	18	Wirth (1972); Versegghy (1974); Krzewicka (2009), Suza, Lich. Bohemoslov. Exs. Fasc. II & V	FL, SR
Bo1	Western Bohemia (Kaiserwald)	Czech Republic	Europe	Cfb	Serpentinite	108	Suza (1927)	FL
Bo2	Křivky National Nature Monument (ex Kaiserwald)	Czech	Europe	Cfb	Serpentinite	127	Peksa (2011)	FL
Bs	Central and SE Bosnia	Bosnia	Europe	Cfb-Cfa	Serpentinite	40	Ritter-Studnička and Klement (1968)	FL
(Bu)	Burgenland	Austria	Europe	Cfb	Serpentinite	3	Hafellner (2001)	SR
Ca	Central coastal range of CA	USA (CA)	N-America	Csb	Serpentinite	68	Sigal (1989)	FL
(ES)	East Sudety	Czech Republic	Europe	Dfb-Cfb	Serpentinite	1	Vězda(1972)	SR
Eu	N-Eubea	Greece	Europe	Csa	Serpentinite	58	Krause and Klement (1962)	FL
Fi	Fidalgo Island	USA (WA)	N-America	Csb	Serpentinite	42	Ryan (1988)	FL
(Ge)	Liguria, Emilia-Romagna, Lombardia (various localities)	Italy	Europe	Csa-Csb	Serpentinite	9	Giordani et al. (2009); Nimis (2016, with refs therein*)	SR
GG	Gurhof-Graben (Melk)	Austria	Europe	Cfb	Serpentinite	35	Kretschmer (1931)	FL
(Gi)	Girvan, Ayrshire	Scotland	Europe	Cfb	Serpentinite	1	Friday (2005)	SR
GV)	Gevne Valley	Turkey	Asia	Csa-Csb	Serpentinite	12	Kocakaya et al. (2014)	FL
GZ	Gostović-Gebiet & Zlatibor-Gebirge; Dobroselica	Serbia-Bosnia	Europe	Cfa-Cfb	Serpentinite	46	Krause and Klement (1958); Poelt (1975)	FL

(IA)	Ile Amsterdam, Ile St. Paul	France	Oceania	Cfb	Serpentinite (exotic)	4	Aptroot et al. (2011)	FL
Ka1	N-Savonia, N-Karelia	Finland	Europe	Dfc	Serpentinite	87	Hakulinen (1958)	FL
Ka2	Niinivaara (N Karelia)	Finland	Europe	Dfc	Serpentinite	116	Takala and Seaward (1978)	FL
KH	Keen of Hamar	Shetland (UK)	Europe	Cfb	Serpentinite	42	Gilbert and James (1987) in; Purvis and Halls (1996)	FL
(Ki)	Mt. Kinabalu, Borneo	Malaysia	Asia	Af-Cfb	Serpentinite	1**	Sipman (1993)	FL
Li	Lizard Peninsula	England	Europe	Cfb	Serpentinite	217	Gilbert and James (1987)	FL
(Ma)	Malatya province	Turkey	Asia	Csa-Dsa	Serpentinite	1	Candan and Turk (2008)	FL
MF	Monte Ferrato	Italy	Europe	Csa-Cfb	Serpentinite	70	Sambo (1927)	FL
Mo	Monviso Massif	Italy	Europe	ET(-Dfc)	Serpentinite	27	Favero-Longo et al. (2015)	FL
(MP)	Monti Pelati	Italy	Europe	Cfa-Cfb	Peridotite, serpentinite	8	Gallo and Piervittori (1991)	FL
Mu	Musiné	Italy	Europe	Cfb-Cfa	Peridotite, serpentinite	29	Favero-Longo et al. (2015)	FL
(NC)	various locations	New Caledonia	Oceania	Aw, Af, Cfa, etc.	Peridotite, serpentinite	14	Aptroot and John (2015)	FL
NI	New Idria	USA (CA)	N-America	Csb	Serpentinite	78	Rajakaruna et al. (2012)	FL
PH	Pine Hill	USA (ME)	N-America	Dfb	Serpentinite	80	Harris et al. (2007), Medeiros et al. (2014)	FL
(Rh)	Rhum	Scotland	Europe	Cfb	Peridotite	19	Gilbert (1983)	FL
SC	Sierra de A Capelada	Spain	Europe	Cfb	Serpentinite	105	Sánchez-Biezma et al. (1996; 2001)	FL; SR
Si	Lower Silesia	Poland	Europe	Cfb	Serpentinite, peridotite	94	Kossowska (2001, with refs therein*)	FL
St	Steiermark	Austria	Europe	Dfc	Serpentinite	99	Hafellner (1991, with refs therein*); Lammermayr (1934)	FL
(SX)	Sierra do Careón	Spain	Europe	Csb	Serpentinite	1	Sánchez-Biezma and López de Silanes (1999)	SR
(TH)	Trial Harbour	Tasmania	Oceania	Cfb	Serpentinite	1	Kantvilas (1991)	SR
(Ti)	Tinos	Greece	Europe	Csa	Serpentinite	17	Zahlbrückner (1907)	FL
Ur	Central Urals	Russia	Asia	Dfb	Serpentinite	116	Paukov and Trapeznikova (2005), Paukov (2009, unpublished data)	FL
(US)	n.a.	USA (PE, MD)	N-America	n.a.	Serpentinite	1	Hansen and Goertzen (2012)	SR
VA	Eastern Valle d'Aosta	Italy	Europe	Cfb-Dfc-ET	Serpentinite	72	Matteucci et al. (2015)	FL
(Vi)	Visegrad	Bosnia	Europe	Cfa-Cfb	Serpentinite	8	Bilovitz and Mayrhofer (2009)	SR

767 *including historical reports; **A list of other 23 species has been recently published for serpentine areas of Borneo (Paukov et al. 2017), which are not considered in total counts
768 and statistical analyses, but reported together with unpublished species reports (Table S7 in Supplementary Materials)

769 Table 2. Areas surveyed with regard to lichen diversity in Valle d'Aosta.
770

Geographic area			Altitude (m a.s.l.)	Lithology	Number of lichen taxa
Abbrev.	Municipality (location)	Coordinates [°]			
GAB	Gressoney-La-Trinité (Gabiét)	N 5079056,E 410423	2340	serpentinite	25
GSB	Saint-Remy-en-Bosses (Grand Saint Bernard)	N 5079930,E 358973	2250	micaschist and gneiss	34
OLL	Ollomont	N 5081308, E 368403	1640	metabasite, chloristoschist	53
SAU	Courmayeur (Pavillon-Saussurea)	N 5077501,E 340665	2200	granite and gneiss	27
THU	La Thuile (Les Granges)	N 5066202,E 341821	1640	micaschist	34
TRI	Gressoney-La-Trinité (village)	N 5075483,E 408645	1680	serpentinite	32
TSA	Saint Cristophe (Tsatelet)	N 5067816,E 370274	570	micaschist	35
VER	Verres (Castle)	N 5058403, E 398430	520	serpentinite	30
VIN	Saint-Vincent	N 5066303, E 395728	650	serpentinite	28

771 [°] one of the four randomly extracted coordinates (UTM ED50) is reported for each area
772

773 Table 3. List of considered functional traits, with the related functional attributes assigned
774 with reference to Smith et al. (2009) and Nimis (2016).

Functional trait	Abbrev.	Functional trait attributes	Abbrev.
Reproductive strategy	REPR	-mainly sexual by means of ascospores -mainly asexual, either by soredia or isidia or fragmentation -with both sexual and asexual structures	Se As Sa
Ascocarp type	ATYP	-with lecanorine apothecia -with lecideine apothecia -with perithecia or others	Ln Ld Pe
Number of ascospores in each ascum	SNUM	-<8 spores -8 spores ->8 spores	<8 =8 >8
Shape of spores	SSHA	-globose, sub-globose, ovate, ellipsoid -different	El Di
Number of septa in the spores	SSEP	-non septate -1 septum ->1 septum	Ns Os Ps
Length of the spores	SLen	-small (<5 µm) -small-medium (5-10 µm) -medium-large (11-20 µm) -large (21-40 µm) -extra-large (>40 µm)	Sl Sm Ml La Xl
Color of the spores	SCOL	-hyaline -from hyaline to dark -dark pigmented	Ia Ch Dp
Growth form	GROW	-crustose -placodioid -squamulose -foliose -fruticose	Cr Cp Sq Fo Fr
Thallus continuity	TCON	-continuous -discontinuous or not visible -with variable continuity	Co Ds De
Photosynthetic strategy	PHOT	-photosynthetic with chlorococcoid green algae -photosynthetic with <i>Trentepohlia</i> pigments -photosynthetic with cyanobacteria pigments	Ch Tr Cy
Color of the thallus	TCOL	-pale -grey -brown-black -orange-yellow -green	Pa Gr Bb Oy Gn
Occurrence of pruina	PRUI	-present -absent -sometimes present	Ye No Yn
Color of the epithecium/disc	ACOL	-black -brown -others	Bl Br Ot
Production of lichen secondary metabolites (categories according to Huneck and Yoshimura 1996):		... for at least one metabolite of a certain category:	
-aliphatic compounds	Alip	-production in all thalli	Yes
-quinones	Quin	-production in some thalli	Y/N
-chromones and xanthones	Chro		
-pulvinic derivatives	Pulv		

-depsides	Deps
-depsidones	Dops
-dibenzofuranes	Dibe
-terpenoids	Terp

775

776

777 Table 4. Percentage contribution from the SDR Simplex analyses of lichen communities in
 778 the ultramafic areas considered at the world scale and ultramafic and non-ultramafic areas
 779 surveyed at the regional scale at different altitudes. S (relative similarity), R (relative
 780 replacement), D (relative richness difference), R+D (relative β -diversity), S+R (relative
 781 richness agreement), S+D (relative nestedness)

	S	R	D	R+D	S+R	S+D	MATRIX FILL (%)
World survey	10.5	55.5	34.0	89.5	66.0	44.3	10.0
Regional survey (ultramafic + non-ultramafic)	24.7	56.1	19.1	75.2	88.8	43.8	9.8
Regional ultramafic	27.6	54.8	17.6	72.4	82.4	45.2	16.4
Regional non- ultramafic	24.9	54.6	20.5	75.0	79.5	45.4	12.0
Regional low alt.	27.4	57.6	15.0	72.6	85.0	42.4	17.9
Regional medium alt.	33.6	45.2	21.2	66.4	78.8	54.8	16.6
Reg high alt.	35.3	43.8	20.9	64.7	79.1	56.2	20.3

783

784 Table 5. Frequency (%) of functional attributes assigned to each trait for species listed in ultramafic areas at the world level (altogether and
 785 separately for groups I-IV of the classification in Fig. 2) and in ultramafic and non-ultramafic areas surveyed at the regional level in Valle d'Aosta
 786 (altogether and separately, for ultramafic, U, and non-ultramafic, NU, areas, and for areas at low, l.alt., medium, m.alt., and high, h.alt., altitude).
 787 Maxima variations of attribute frequencies among groups I-IV at the world scale (ΔGr) and among different substrates (ΔU -NU) and altitudes (Δalt)
 788 at the regional scale (underlined values indicate a higher maximum frequency variation among substrates or altitudes for a certain attribute).
 789 Abbreviations for functional traits and attributes are listed in Table 3.

Functional traits	Functional attributes	World						Regional							
		Overall	Gr. I	Gr. II	Gr. III	Gr. IV	ΔGr (max)	Overall	U	NU	l.alt.	m.alt.	h.alt.	ΔU -NU	Δalt (max)
REPR	Se	65.2	64.6	63.7	69.6	76.2	12.5	86.2	88.7	86.0	86.8	85.3	90.2	2.7	<u>4.9</u>
	As	6.1	2.5	6.5	3.8	4.8	3.9	0.9	0.0	1.1	0.0	1.3	0.0	1.1	<u>1.3</u>
	Sa	28.8	32.9	29.8	26.6	19.0	13.9	12.8	11.3	12.9	13.2	13.3	9.8	1.6	<u>3.5</u>
ATYP	Ln	46.2	49.4	46.8	41.8	47.6	7.6	33.0	30.6	31.2	24.5	29.3	47.1	0.5	<u>22.5</u>
	Ld	50.0	45.6	50.0	54.4	47.6	8.9	59.6	61.3	61.3	73.6	61.3	45.1	0.0	<u>28.5</u>
	Pe	3.8	5.1	3.2	3.8	4.8	1.8	7.3	8.1	7.5	1.9	9.3	7.8	0.5	<u>7.4</u>
SNUM	<8	6.8	3.8	7.3	7.6	6.0	3.8	7.3	9.7	7.5	11.3	9.3	7.8	2.2	<u>3.5</u>
	8.0	90.2	91.1	90.3	89.9	89.3	1.0	81.7	83.9	79.6	77.4	78.7	80.4	<u>4.3</u>	3.0
	>8	3.0	5.1	2.4	2.5	4.8	2.6	11.0	6.5	12.9	11.3	12.0	11.8	<u>6.5</u>	0.7
SSHA	El	93.2	94.9	92.7	94.9	97.6	4.9	94.5	96.8	94.6	94.3	96.0	96.1	<u>2.2</u>	1.7
	Di	6.8	5.1	7.3	5.1	2.4	4.9	5.5	3.2	5.4	5.7	4.0	3.9	<u>2.2</u>	1.7
	Ns	73.5	72.2	75.0	73.4	70.2	4.8	68.8	64.5	74.2	62.3	70.7	76.5	9.7	<u>14.2</u>
SSEP	Os	13.6	16.5	12.9	13.9	17.9	5.0	26.6	29.0	21.5	34.0	22.7	19.6	7.5	<u>14.4</u>
	Ps	12.9	11.4	12.1	12.7	11.9	1.3	4.6	6.5	4.3	3.8	6.7	3.9	2.2	<u>2.9</u>
	Sl	3.8	2.5	4.0	2.5	2.4	1.7	9.2	6.5	10.8	11.3	9.3	11.8	<u>4.3</u>	2.4
SLEN	Sm	11.4	12.7	12.1	7.6	11.9	5.1	22.0	21.0	21.5	22.6	18.7	13.7	0.5	<u>8.9</u>
	MI	63.6	60.8	64.5	69.6	64.3	8.9	52.3	58.1	50.5	49.1	54.7	56.9	7.5	<u>7.8</u>
	La	14.4	19.0	13.7	15.2	17.9	5.3	13.8	11.3	15.1	13.2	13.3	15.7	<u>3.8</u>	2.5
SCOL	XI	6.8	5.1	5.6	5.1	3.6	2.1	2.8	3.2	2.2	3.8	4.0	2.0	1.1	<u>2.0</u>
	Ia	86.4	86.1	87.1	86.1	83.3	3.8	83.5	80.6	86.0	81.1	82.7	86.3	<u>5.4</u>	5.1
	Ch	1.5	0.0	1.6	0.0	0.0	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
GROW	Dp	12.1	13.9	11.3	13.9	16.7	5.4	16.5	19.4	14.0	18.9	17.3	13.7	<u>5.4</u>	5.1
	Cr	56.1	59.5	53.2	55.7	65.5	12.3	67.0	67.7	64.5	67.9	64.0	78.4	3.2	<u>14.4</u>
	Cp	2.3	1.3	2.4	2.5	3.6	2.3	2.8	3.2	3.2	5.7	2.7	2.0	0.0	<u>3.7</u>

	Sq	2.3	2.5	2.4	0.0	3.6	3.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Fo	15.9	20.3	16.9	16.5	14.3	6.0	26.6	25.8	28.0	26.4	29.3	13.7	2.2	<u>15.6</u>
	Fr	23.5	16.5	25.0	25.3	13.1	12.2	3.7	3.2	4.3	0.0	4.0	5.9	1.1	<u>5.9</u>
TCON	Co	83.3	79.7	82.3	84.8	78.6	6.2	78.0	82.3	79.6	79.2	77.3	78.4	2.7	1.9
	Ds	1.5	2.5	1.6	0.0	2.4	2.5	5.5	6.5	4.3	9.4	5.3	2.0	2.2	<u>7.5</u>
	De	15.2	17.7	16.1	15.2	19.0	3.9	16.5	11.3	16.1	11.3	17.3	19.6	4.8	<u>8.3</u>
PHOT	Ch	90.2	89.9	89.5	93.7	92.9	4.2	100.0	100.0	100.0	100.0	100.0	100.0	0.0	0.0
	Tr	0.8	1.3	0.8	1.3	1.2	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Cy	9.1	8.9	9.7	5.1	6.0	4.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TCOL	Pa	22.0	20.3	19.4	22.8	28.6	9.2	28.4	19.4	30.1	22.6	28.0	31.4	<u>10.8</u>	8.7
	Gr	35.6	40.5	36.3	36.7	34.5	6.0	33.9	41.9	31.2	34.0	36.0	37.3	<u>10.8</u>	3.3
	Bb	18.9	20.3	20.2	12.7	16.7	7.6	20.2	16.1	23.7	13.2	24.0	23.5	7.5	<u>10.8</u>
	Oy	6.1	8.9	5.6	8.9	6.0	3.2	7.3	6.5	7.5	9.4	6.7	5.9	1.1	<u>3.6</u>
	Gn	17.4	10.1	18.5	19.0	14.3	8.9	10.1	16.1	7.5	20.8	5.3	2.0	8.6	<u>18.8</u>
PRUI	Ye	12.9	10.1	13.7	12.7	16.7	6.5	27.5	33.9	25.8	35.8	25.3	15.7	8.1	<u>20.2</u>
	Yn	4.5	6.3	4.8	5.1	3.6	2.8	1.8	1.6	2.2	0.0	2.7	2.0	0.5	<u>2.7</u>
	No	82.6	83.5	81.5	82.3	79.8	3.8	70.6	64.5	72.0	64.2	72.0	82.4	7.5	<u>18.2</u>
ACOL	Bl	35.6	38.0	33.1	39.2	47.6	14.6	56.9	53.2	59.1	43.4	58.7	72.5	5.9	<u>29.2</u>
	Br	46.2	46.8	48.4	41.8	34.5	13.9	29.4	32.3	29.0	32.1	29.3	19.6	3.2	<u>12.5</u>
	Ot	18.2	15.2	18.5	19.0	17.9	3.8	13.8	14.5	11.8	24.5	12.0	7.8	2.7	<u>16.7</u>
Alip	Yes	8.3	3.8	8.9	7.6	10.7	6.9	6.4	9.7	7.5	11.3	8.0	13.7	2.2	<u>5.7</u>
	Y/N	1.5	1.3	1.6	1.3	1.2	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Quin	Yes	4.5	5.1	4.0	5.1	4.8	1.0	8.3	9.7	5.4	13.2	6.7	2.0	4.3	<u>11.2</u>
	Y/N	0.8	1.3	0.8	0.0	0.0	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Chro	Yes	3.8	3.8	4.0	3.8	4.8	0.2	3.7	1.6	4.3	3.8	2.7	5.9	2.7	<u>3.2</u>
	Y/N	0.8	1.3	0.8	0.0	1.2	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pulv	Yes	3.0	5.1	3.2	5.1	4.8	1.8	3.7	3.2	4.3	5.7	4.0	3.9	1.1	<u>1.7</u>
	Y/N	4.8	2.5	1.6	2.5	2.4	0.9	0.9	1.6	0.0	0.0	1.3	0.0	<u>1.6</u>	1.3
Deps	Yes	35.6	36.7	36.3	34.2	31.0	5.8	29.4	25.8	31.2	20.8	30.7	33.3	<u>5.4</u>	2.7
	Y/N	6.8	7.6	6.5	10.1	4.8	5.4	3.7	4.8	4.3	3.8	5.3	5.9	0.5	<u>2.1</u>
Dops	Yes	27.3	24.1	24.2	34.2	27.4	10.1	29.4	37.1	30.1	35.8	30.7	21.6	7.0	<u>14.3</u>
	Y/N	11.4	11.4	11.3	11.4	13.1	1.8	14.7	17.7	14.0	11.3	16.0	21.6	3.8	<u>10.2</u>
Dibe	Yes	8.3	2.5	8.9	7.6	9.5	7.0	4.6	6.5	5.4	9.4	5.3	3.9	1.1	<u>5.5</u>
	Y/N	1.5	2.5	1.6	2.5	0.0	2.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Terp	Yes	7.6	7.6	8.1	5.1	9.5	4.5	7.3	4.8	8.6	7.5	8.0	9.8	<u>3.8</u>	2.3
	Y/N	3.8	1.3	4.0	2.5	1.2	2.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

790 **Supplementary materials**

791 Table S1 Checklist of lichen taxa reported for serpentine areas worldwide. Abbreviation of
792 geographic areas are reported in Table 1.

793 (attached Excel file)

794 Table S2 Functional traits of lichen taxa occurring in at least five ultramafic areas worldwide
795 (i.e. species reported in >20% of sites with at least 20 species).

796 (attached Excel file)

797 Table S3 Scores of CCA-I.

798 (attached Excel file)

799 Table S4 List of lichen taxa surveyed in the surveyed serpentine and non-serpentine areas of
800 Aosta Valley. Abbreviation of sites are reported in Table 2.

801 (attached Excel file)

802 Table S5 Functional traits of lichen taxa surveyed at the regional level in ultramafic and non-
803 ultramafic areas of Valle d'Aosta (NW-Italy).

804 (attached Excel file)

805 Table S6 Scores of CCA-II.

806 (attached Excel file)

807 Table S7 Recently published and work-in-progress information on lichen diversity in
808 serpentine areas.

809 (attached Excel file)

810 Table S8 Thirteen species and varieties reported as ultramafic endemic lichens, i.e. growing
811 exclusively on serpentinite and related rocks (Favero-Longo et al. 2004).

812 (attached Word file)

Figure 1




Figure 2



Figure 3




Figure 4

[Click here to download Figure Fig.4.tif](#)




Figure 5




Figure 6

[Click here to download Figure Fig.6.tif](#)





Figure 7

[Click here to download Figure Fig.7.tif](#)





[Click here to access/download](#)

Electronic Supplementary Material
Favero-Longo_et_al_Table S8.pdf





[Click here to access/download](#)

Electronic Supplementary Material

Favero-Longo_et_al_TableS1-S7 unmod_bis.xlsx

